


Pooled Association Tests for Rare Variants in Exon-Resequencing Studies

Alkes L. Price, Gregory V. Kryukov, Paul I. W. de Bakker, Shaun M. Purcell, Jeff Staples, Lee-Jen Wei, and Shamil R. Sunyaev


Figure S1. Prevalence of Goldilocks Alleles in Simulations at Different Values of Selection

Parameter s

(a)



(b)



We consider $s=0.01$ ($\log_{10}s=-2$), $s=0.001$ ($\log_{10}s=-3$), or $s=0.0001$ ($\log_{10}s=-4$). We plot (a) the fraction of all functional SNPs that have frequency between 0.5% and 2%, and (b) the probability that a gene has at least one functional SNP with frequency between 0.5% and 2%, as a function of s .

Table S1. Power of Various Approaches Using Quantitative Phenotypes

	T1	T5	WE	VT
$s=0.0001, \delta=0.125, \alpha=0.001$	0.029	0.082	0.030	0.108
$s=0.0001, \delta=0.125, \alpha=0.05$	0.298	0.386	0.313	0.431
$s=0.0001, \delta=0.25, \alpha=0.001$	0.228	0.369	0.243	0.487
$s=0.0001, \delta=0.25, \alpha=0.05$	0.672	0.706	0.734	0.805
$s=0.0001, \delta=0.5, \alpha=0.001$	0.788	0.777	0.890	0.925
$s=0.0001, \delta=0.5, \alpha=0.05$	0.954	0.925	0.983	0.985
$s=0.001, \delta=0.125, \alpha=0.001$	0.015	0.034	0.013	0.030
$s=0.001, \delta=0.125, \alpha=0.05$	0.230	0.249	0.220	0.249
$s=0.001, \delta=0.25, \alpha=0.001$	0.135	0.180	0.097	0.205
$s=0.001, \delta=0.25, \alpha=0.05$	0.547	0.502	0.545	0.598
$s=0.001, \delta=0.5, \alpha=0.001$	0.569	0.492	0.596	0.693
$s=0.001, \delta=0.5, \alpha=0.05$	0.892	0.792	0.941	0.947
$s=0.01, \delta=0.125, \alpha=0.001$	0.001	0.001	0.002	0.001
$s=0.01, \delta=0.125, \alpha=0.05$	0.060	0.056	0.066	0.066
$s=0.01, \delta=0.25, \alpha=0.001$	0.002	0.002	0.002	0.003
$s=0.01, \delta=0.25, \alpha=0.05$	0.075	0.068	0.082	0.082
$s=0.01, \delta=0.5, \alpha=0.001$	0.004	0.003	0.006	0.006
$s=0.01, \delta=0.5, \alpha=0.05$	0.111	0.090	0.132	0.129

We display results for T1 (1% allele frequency threshold), T5 (5% allele frequency threshold), WE (weighted) and VT (variable threshold) analyses for various values of s and δ . Results are displayed for $\alpha=0.001$ and $\alpha=0.05$ significance levels, based on 10,000 independent simulations. The best-performing method(s) are indicated in bold. We note that of 54 pairwise comparisons between VT and one other

method, VT performs worse in 3/54 comparisons, with the reduction in power equal to 0.004 or less in each case.

Table S2

	T1	T5	WE	VT
$\alpha=0.001$	0.098	0.223	0.153	0.400
$\alpha=0.05$	0.406	0.503	0.517	0.630

(a) Power of various approaches using quantitative phenotypes, for a simulation in which phenotypes are independent of selection coefficient. The fraction of functional mutations, i.e. mutations that affect phenotype, was equal to the fraction of mutations with selection coefficient above 10^{-3} (48%), however the functionality of each mutation was independent of the selection coefficient. We display results for T1 (1% allele frequency threshold), T5 (5% allele frequency threshold), WE (weighted), and VT (variable threshold) analyses for $\alpha=0.001$ and $\alpha=0.05$ significance levels, based on 10,000 independent simulations.\

	T1	T5	WE	VT	VTP	VTP-hi	VTP-lo
$\alpha=0.001$	0.135	0.180	0.097	0.205	0.257	0.269	0.231
$\alpha=0.05$	0.547	0.502	0.545	0.598	0.686	0.698	0.640

(b) Value of incorporation of PolyPhen-2 predictions, as a function of the (known) distribution of PolyPhen-2 scores associated to either deep or shallow multiple sequence alignments. Alignments with less than 10 sequences (after PolyPhen-2 filtering for very close homologs) were classified as “shallow”. Alignments with 100 or more sequences were classified as “deep”. We display results for T1 (1% allele frequency threshold), T5 (5% allele frequency threshold), WE (weighted), VT (variable threshold), VTP (VT plus PolyPhen-2), VTP-deep (VT plus PolyPhen-2, deep alignments) and VTP-shallow (VT plus PolyPhen-2, shallow alignments) analyses for $\alpha=0.001$ and $\alpha=0.05$ significance levels, based on 10,000

independent simulations. All columns except VTP-deep and VTP-shallow are identical to Table 1. As expected, VTP-deep outperforms VTP-shallow, but even VTP-shallow is substantially better than VT.

Table S3. Power of Various Approaches Using Dichotomous Phenotypes

	T1	T5	WE	VT
$s=0.0001, \delta=0.125, \alpha=0.001$	0.022	0.060	0.029	0.088
$s=0.0001, \delta=0.125, \alpha=0.05$	0.258	0.340	0.302	0.398
$s=0.0001, \delta=0.25, \alpha=0.001$	0.153	0.306	0.222	0.435
$s=0.0001, \delta=0.25, \alpha=0.05$	0.608	0.659	0.704	0.771
$s=0.0001, \delta=0.5, \alpha=0.001$	0.690	0.709	0.836	0.887
$s=0.0001, \delta=0.5, \alpha=0.05$	0.930	0.903	0.984	0.985
$s=0.001, \delta=0.125, \alpha=0.001$	0.012	0.023	0.011	0.021
$s=0.001, \delta=0.125, \alpha=0.05$	0.202	0.219	0.205	0.233
$s=0.001, \delta=0.25, \alpha=0.001$	0.089	0.141	0.079	0.163
$s=0.001, \delta=0.25, \alpha=0.05$	0.474	0.451	0.487	0.528
$s=0.001, \delta=0.5, \alpha=0.001$	0.438	0.410	0.480	0.578
$s=0.001, \delta=0.5, \alpha=0.05$	0.838	0.737	0.891	0.901
$s=0.01, \delta=0.125, \alpha=0.001$	0.001	0.001	0.001	0.002
$s=0.01, \delta=0.125, \alpha=0.05$	0.060	0.057	0.062	0.064
$s=0.01, \delta=0.25, \alpha=0.001$	0.002	0.002	0.002	0.002
$s=0.01, \delta=0.25, \alpha=0.05$	0.070	0.064	0.076	0.076
$s=0.01, \delta=0.5, \alpha=0.001$	0.003	0.003	0.004	0.004
$s=0.01, \delta=0.5, \alpha=0.05$	0.095	0.080	0.111	0.107

We display results for T1 (1% allele frequency threshold), T5 (5% allele frequency threshold), WE (weighted) and VT (variable threshold) analyses for various values of s and δ . Results are displayed for for $\alpha=0.001$ and $\alpha=0.05$ significance levels, based on 10,000 independent simulations. The best-performing method(s) are indicated in bold. We note that of 54 pairwise comparisons between VT and

one other method, VT performs worse in 2/54 comparisons, with the reduction in power equal to 0.004 or less in each case.

Table S4

	T1		T5		WE		VT	
	-PolyPhen	+PolyPhen	-PolyPhen	+PolyPhen	-PolyPhen	+PolyPhen	-PolyPhen	+PolyPhen
$\alpha=0.001$	0.14	0.22	0.18	0.24	0.10	0.15	0.21	0.26
$\alpha=0.05$	0.55	0.69	0.50	0.62	0.55	0.65	0.60	0.69

(a) Effect of PolyPhen-2 on power of various approaches using quantitative phenotypes. We display results for T1 (1% allele frequency threshold), T5 (5% allele frequency threshold), WE (weighted), and VT (variable threshold) analyses (each with and without PolyPhen-2) for $\alpha=0.001$ and $\alpha=0.05$ significance levels, based on 10,000 independent simulations.

	T1		T5		WE		VT	
	-PolyPhen	+PolyPhen	-PolyPhen	+PolyPhen	-PolyPhen	+PolyPhen	-PolyPhen	+PolyPhen
$\alpha=0.001$	0.09	0.15	0.14	0.19	0.08	0.12	0.16	0.21
$\alpha=0.05$	0.47	0.62	0.45	0.56	0.49	0.60	0.53	0.62

(b) Effect of PolyPhen-2 on power of various approaches using dichotomous phenotypes. We display results for T1 (1% allele frequency threshold), T5 (5% allele frequency threshold), WE (weighted), and VT (variable threshold) analyses (each with and without PolyPhen-2) for $\alpha=0.001$ and $\alpha=0.05$ significance levels, based on 10,000 independent simulations.